

【講座 精神疾患の臨床】1 気分症群  
p229-239 「双極症の生物学的研究」(加藤忠史)

参考文献

- 1) Akula N, et al. RNA-sequencing of the brain transcriptome implicates dysregulation of neuroplasticity, circadian rhythms and GTPase binding in bipolar disorder. *Mol Psychiatry* 2014; 19 (11): 1179–1185.
- 2) Altshuler L, et al. Increased amygdala activation during mania: A functional magnetic resonance imaging study. *Am J Psychiatry* 2005; 162 (6): 1211–1213.
- 3) Bamne MN, et al. Application of an ex vivo cellular model of circadian variation for bipolar disorder research: A proof of concept study. *Bipolar Disord* 2013; 15 (6): 694–700.
- 4) Baumann B, et al. Tyrosine hydroxylase immunoreactivity in the locus coeruleus is reduced in depressed non-suicidal patients but normal in depressed suicide patients. *Eur Arch Psychiatry Clin Neurosci* 1999; 249 (4): 212–219.
- 5) Baumann B, et al. Circumscribed numerical deficit of dorsal raphe neurons in mood disorders. *Psychol Med* 2002; 32 (1): 93–103.
- 6) Bavamian S, et al. Dysregulation of miR-34a links neuronal development to genetic risk factors for bipolar disorder. *Mol Psychiatry* 2015; 20 (5): 573–584.
- 7) Belmaker RH, et al. Reduced inositol content in lymphocyte-derived cell lines from bipolar patients. *Bipolar Disord* 2002; 4 (1): 67–69.
- 8) Benes FM, et al. A reduction of nonpyramidal cells in sector CA2 of schizophrenics and manic depressives. *Biol Psychiatry* 1998; 44 (2): 88–97.
- 9) Benes FM, et al. Regulation of the GABA cell phenotype in hippocampus of schizophrenics and bipolar patients. *Proc Natl Acad Sci U S A* 2007; 104 (24): 10164–10169.
- 10) Bestelmeyer PE, et al. The P300 as a possible endophenotype for schizophrenia and bipolar disorder: Evidence from twin and patient studies. *Psychiatry Res* 2009; 169 (3): 212–219.
- 11) Bielau H, et al. Volume deficits of subcortical nuclei in mood disorders: A postmortem study. *Eur Arch Psychiatry Clin Neurosci* 2005; 255 (6): 401–412.
- 12) Blumberg HP, et al. A functional magnetic resonance imaging study of bipolar disorder: State- and trait-related dysfunction in ventral prefrontal cortices. *Arch Gen Psychiatry* 2003; 60 (6): 601–609.
- 13) Blumberg HP, et al. Preliminary evidence for medication effects on functional abnormalities in the amygdala and anterior cingulate in bipolar disorder. *Psychopharmacology (Berl)* 2005; 183 (3): 308–313.
- 14) Breen MS, et al. Lithium-responsive genes and gene networks in bipolar disorder patient-derived lymphoblastoid cell lines. *Pharmacogenomics J* 2016; 16 (5): 446–453.
- 15) Breen MS, et al. Dynamic landscape and genetic regulation of RNA editing in schizophrenia. *bioRxiv* 2019.
- 16) Cannon DM, et al. Serotonin transporter binding in bipolar disorder assessed using [11C]DASB and positron emission tomography. *Biol Psychiatry* 2006; 60 (3): 207–217.
- 17) Cannon DM, et al. Elevated serotonin transporter binding in major depressive disorder assessed using positron emission tomography and [11C]DASB; comparison with bipolar disorder. *Biol Psychiatry* 2007; 62 (8): 870–877.
- 18) Chen C, et al. Correlation between DNA methylation and gene expression in the brains of patients with bipolar disorder and schizophrenia. *Bipolar Disord* 2014; 16 (8): 790–799.
- 19) Chen HM, et al. Transcripts involved in calcium signaling and telencephalic neuronal fate are altered in induced pluripotent stem cells from bipolar disorder patients. *Transl Psychiatry* 2014; 4: e375.
- 20) Chuang DM, et al. GSK-3 as a target for lithium-induced neuroprotection against excitotoxicity in neuronal cultures and animal models of ischemic stroke. *Front Mol Neurosci* 2011; 4: 15.
- 21) Cotter D, et al. Reduced glial cell density and neuronal size in the anterior cingulate cortex in major depressive disorder. *Arch Gen Psychiatry* 2001; 58 (6): 545–553.
- 22) Coufal NG, et al. L1 retrotransposition in human neural progenitor cells. *Nature* 2009; 460 (7259): 1127–1131.
- 23) Craddock N, et al. Familial cosegregation of major affective disorder and Darier's disease (Keratosis Follicularis). *Br J Psychiatry* 1994; 164: 355–358.

- 24) Cruz DA, et al. Selective alterations in postsynaptic markers of chandelier cell inputs to cortical pyramidal neurons in subjects with schizophrenia. *Neuropsychopharmacology* 2009; 34 (9): 2112–2124.
- 25) Curtis VA, et al. Differential frontal activation in schizophrenia and bipolar illness during verbal fluency. *J Affect Disord* 2001; 66 (2–3): 111–121.
- 26) Dao DT, et al. Mood disorder susceptibility gene CACNA1C modifies mood-related behaviors in mice and interacts with sex to influence behavior in mice and diagnosis in humans. *Biol Psychiatry* 2010; 68 (9): 801–810.
- 27) Dogan AE, et al. Brain lactate and pH in schizophrenia and bipolar disorder: A systematic review of findings from magnetic resonance studies. *Neuropsychopharmacology* 2018; 43 (8): 1681–1690.
- 28) Dokucu ME, et al. Lithium- and valproate-induced alterations in circadian locomotor behavior in Drosophila. *Neuropsychopharmacology* 2005; 30 (12): 2216–2224.
- 29) Dowlatshahi D, et al. Increased hippocampal supragranular Timm staining in subjects with bipolar disorder. *Neuroreport* 2000; 11 (17): 3775–3778.
- 30) Doyle GA, et al. Analysis of LINE-1 elements in DNA from postmortem brains of individuals with schizophrenia. *Neuropsychopharmacology* 2017; 42 (13): 2602–2611.
- 31) Dracheva S, et al. Increased serotonin 2C receptor mRNA editing: A possible risk factor for suicide. *Mol Psychiatry* 2008; 13 (11): 1001–1010.
- 32) Etain B, et al. Genetic and functional abnormalities of the melatonin biosynthesis pathway in patients with bipolar disorder. *Hum Mol Genet* 2012; 21 (18): 4030–4037.
- 33) Fattal O, et al. Psychiatric comorbidity in 36 adults with mitochondrial cytopathies. *CNS Spectr* 2007; 12 (6): 429–438.
- 34) Ferreira MA, et al. Collaborative genome-wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. *Nat Genet* 2008; 40 (9): 1056–1058.
- 35) Föcking M, et al. Common proteomic changes in the hippocampus in schizophrenia and bipolar disorder and particular evidence for involvement of cornu ammonis regions 2 and 3. *Arch Gen Psychiatry* 2011; 68 (5): 477–488.
- 36) Frank E, et al. Two-year outcomes for interpersonal and social rhythm therapy in individuals with bipolar I disorder. *Arch Gen Psychiatry* 2005; 62 (9): 996–1004.
- 37) Frey BN, et al. Increased oxidative stress and DNA damage in bipolar disorder: A twin-case report. *Prog Neuropsychopharmacol Biol Psychiatry* 2007; 31 (1): 283–285.
- 38) Gaspar L, et al. Human cellular differences in cAMP--CREB signaling correlate with light-dependent melatonin suppression and bipolar disorder. *Eur J Neurosci* 2014; 40 (1): 2206–2215.
- 39) Gruber SA, et al. Decreased activation of the anterior cingulate in bipolar patients: An fMRI study. *J Affect Disord* 2004; 82 (2): 191–201.
- 40) Haarman BC, et al. Neuroinflammation in bipolar disorder - A [(11)C]-(R)-PK11195 positron emission tomography study. *Brain Behav Immun* 2014; 40: 219–225.
- 41) Hahn CG, et al. Aberrant intracellular calcium signaling in olfactory neurons from patients with bipolar disorder. *Am J Psychiatry* 2005; 162 (3): 616–618.
- 42) Harwood AJ. Lithium and bipolar mood disorder: The inositol-depletion hypothesis revisited. *Mol Psychiatry* 2005; 10 (1): 117–126.
- 43) Hayashi A, et al. Aberrant endoplasmic reticulum stress response in lymphoblastoid cells from patients with bipolar disorder. *Int J Neuropsychopharmacol* 2009; 12 (1): 33–43.
- 44) Huang JH, et al. Perturbational profiling of metabolites in patient fibroblasts implicates alpha-amino adipate as a potential biomarker for bipolar disorder. *Mol Neuropsychiatry* 2016; 2 (2): 97–106.
- 45) Ichimiya T, et al. Serotonin transporter binding in patients with mood disorders: A PET study with [11C](+)-McN5652. *Biol Psychiatry* 2002; 51 (9): 715–722.
- 46) Inczedy-Farkas G, et al. Psychiatric symptoms of patients with primary mitochondrial DNA disorders. *Behav Brain Funct* 2012; 8: 9.
- 47) Iwamoto K, Kato T. RNA editing of serotonin 2C receptor in human postmortem brains of major mental disorders. *Neurosci Lett* 2003; 346 (3): 169–172.
- 48) Iwamoto K, et al. Molecular characterization of bipolar disorder by comparing gene expression profiles of postmortem brains of major mental disorders. *Mol Psychiatry* 2004; 9 (4): 406–416.
- 49) Iwamoto K, et al. Altered expression of mitochondria-related genes in postmortem brains of patients with bipolar disorder or schizophrenia, as revealed by large-scale DNA microarray analysis. *Hum Mol*

- Genet* 2005; 14 (2): 241–253.
- 50) Iwamoto K, et al. Neurons show distinctive DNA methylation profile and higher interindividual variations compared with non-neurons. *Genome Res* 2011; 21 (5): 688–696.
  - 51) Kageyama Y, et al. The relationship between circulating mitochondrial DNA and inflammatory cytokines in patients with major depression. *J Affect Disord* 2018; 233: 15–20.
  - 52) Kam JW, et al. Resting state EEG power and coherence abnormalities in bipolar disorder and schizophrenia. *J Psychiatr Res* 2013; 47 (12): 1893–1901.
  - 53) Kaminsky Z, et al. A multi-tissue analysis identifies HLA complex group 9 gene methylation differences in bipolar disorder. *Mol Psychiatry* 2012; 17 (7): 728–740.
  - 54) Kasahara T, et al. Mice with neuron-specific accumulation of mitochondrial DNA mutations show mood disorder-like phenotypes. *Mol Psychiatry* 2006; 11 (6): 577–593.
  - 55) Kato T, et al. Increased levels of a mitochondrial DNA deletion in the brain of patients with bipolar disorder. *Biol Psychiatry* 1997; 42 (10): 871–875.
  - 56) Kato T, et al. Behavioral and gene expression analyses of Wfs1 knockout mice as a possible animal model of mood disorder. *Neurosci Res* 2008; 61 (2): 143–158.
  - 57) Kato T, et al. Gene expression analysis in lymphoblastoid cells as a potential biomarker of bipolar disorder. *J Hum Genet* 2011; 56 (11): 779–783.
  - 58) Kim DJ, et al. Disturbed resting state EEG synchronization in bipolar disorder: A graph-theoretic analysis. *Neuroimage Clin* 2013; 2: 414–423.
  - 59) Kim KH, et al. Transcriptomic analysis of induced pluripotent stem cells derived from patients with bipolar disorder from an old order amish pedigree. *PLoS One* 2015; 10 (11): e0142693.
  - 60) Kimmel RJ, et al. Cosegregation of bipolar disorder and autosomal-dominant medullary cystic kidney disease in a large family. *Am J Psychiatry* 2005; 162 (10): 1972–1974.
  - 61) King DP, et al. Positional cloning of the mouse circadian clock gene. *Cell* 1997; 89 (4): 641–653.
  - 62) Klein PS, Melton DA. A molecular mechanism for the effect of lithium on development. *Proc Natl Acad Sci U S A* 1996; 93 (16): 8455–8459.
  - 63) Kohen R, et al. Transcriptome profiling of human hippocampus dentate gyrus granule cells in mental illness. *Transl Psychiatry* 2014; 4: e366.
  - 64) Konopaske GT, et al. Prefrontal cortical dendritic spine pathology in schizophrenia and bipolar disorder. *JAMA Psychiatry* 2014; 71 (12): 1323–1331.
  - 65) Konradi C, et al. Molecular evidence for mitochondrial dysfunction in bipolar disorder. *Arch Gen Psychiatry* 2004; 61 (3): 300–308.
  - 66) Konradi C, et al. Hippocampal interneurons in bipolar disorder. *Arch Gen Psychiatry* 2011; 68 (4): 340–350.
  - 67) Kubota M, et al. Abnormal Ca<sup>2+</sup> dynamics in transgenic mice with neuron-specific mitochondrial DNA defects. *J Neurosci* 2006; 26 (47): 12314–12324.
  - 68) Kubota-Sakashita M, et al. A role of ADAR2 and RNA editing of glutamate receptors in mood disorders and schizophrenia. *Mol Brain* 2014; 7: 5.
  - 69) Kuloglu M, et al. Antioxidant enzyme activities and malondialdehyde levels in patients with obsessive-compulsive disorder. *Neuropsychobiology* 2002; 46 (1): 27–32.
  - 70) Lavebratt C, et al. The KMO allele encoding Arg452 is associated with psychotic features in bipolar disorder type 1, and with increased CSF KYNA level and reduced KMO expression. *Mol Psychiatry* 2014; 19 (3): 334–341.
  - 71) Lee PS, et al. Distinct neuronal oscillatory responses between patients with bipolar and unipolar disorders: A magnetoencephalographic study. *J Affect Disord* 2010; 123 (1–3): 270–275.
  - 72) Lewy AJ, et al. Manic-depressive patients may be supersensitive to light. *Lancet (London, England)* 1981; 1 (8216): 383–384.
  - 73) Lyddon R, et al. Ionotropic glutamate receptor mRNA editing in the prefrontal cortex: No alterations in schizophrenia or bipolar disorder. *J Psychiatry Neurosci* 2012; 37 (4): 267–272.
  - 74) Machado-Vieira R, et al. Decreased plasma brain derived neurotrophic factor levels in unmedicated bipolar patients during manic episode. *Biol Psychiatry* 2007; 61 (2): 142–144.
  - 75) Malhi GS, et al. An emotional Stroop functional MRI study of euthymic bipolar disorder. *Bipolar Disord* 2005; 7 (Suppl 5): 58–69.
  - 76) Manaye KF, et al. Selective neuron loss in the paraventricular nucleus of hypothalamus in patients suf-

- fering from major depression and bipolar disorder. *J Neuropathol Exp Neurol* 2005; 64 (3): 224–229.
- 77) Mancuso M, et al. Psychiatric involvement in adult patients with mitochondrial disease. *Neurol Sci* 2013; 34 (1): 71–74.
- 78) Mansour HA, et al. Circadian phase variation in bipolar I disorder. *Chronobiol Int* 2005; 22 (3): 571–584.
- 79) Matthews PR, Harrison PJ. A morphometric, immunohistochemical, and in situ hybridization study of the dorsal raphe nucleus in major depression, bipolar disorder, schizophrenia, and suicide. *J Affect Disord* 2012; 137 (1–3): 125–134.
- 80) McCarthy MJ, et al. Genetic and clinical factors predict lithium's effects on PER2 gene expression rhythms in cells from bipolar disorder patients. *Transl Psychiatry* 2013; 3: e318.
- 81) McCarthy MJ, et al. Disinhibition of the extracellular-signal-regulated kinase restores the amplification of circadian rhythms by lithium in cells from bipolar disorder patients. *Eur Neuropsychopharmacol* 2016; 26 (8): 1310–1319.
- 82) McClung CA, et al. Regulation of dopaminergic transmission and cocaine reward by the Clock gene. *Proc Natl Acad Sci U S A* 2005; 102 (26): 9377–9381.
- 83) Meyer U, Feldon J. Neural basis of psychosis-related behaviour in the infection model of schizophrenia. *Behav Brain Res* 2009; 204 (2): 322–334.
- 84) Mill J, et al. Epigenomic profiling reveals DNA-methylation changes associated with major psychosis. *Am J Hum Genet* 2008; 82 (3): 696–711.
- 85) Miller JM, et al. Positron emission tomography quantification of serotonin transporter binding in medication-free bipolar disorder. *Synapse* 2016; 70 (1): 24–32.
- 86) Morsel AM, et al. Systematic review of cognitive event related potentials in euthymic bipolar disorder. *Clin Neurophysiol* 2018; 129 (9): 1854–1865.
- 87) Mukherjee S, et al. Knockdown of Clock in the ventral tegmental area through RNA interference results in a mixed state of mania and depression-like behavior. *Biol Psychiatry* 2010; 68 (6): 503–511.
- 88) Nanko S, et al. Organic mood syndrome in two siblings with Wolfram syndrome. *Br J Psychiatry* 1992; 161: 282.
- 89) Nemanov L, et al. Effect of bipolar disorder on lymphocyte inositol monophosphatase mRNA levels. *Int J Neuropsychopharmacol* 1999; 2 (1): 25–29.
- 90) Nishioka M, et al. Identification of somatic mutations in postmortem human brains by whole genome sequencing and their implications for psychiatric disorders. *Psychiatry Clin Neurosci* 2018; 72 (4): 280–294.
- 91) Nurberger JL, Jr., et al. Melatonin suppression by light in euthymic bipolar and unipolar patients. *Arch Gen Psychiatry* 2000; 57 (6): 572–579.
- 92) O'Donnell T, et al. Chronic lithium and sodium valproate both decrease the concentration of myoinositol and increase the concentration of inositol monophosphates in rat brain. *Eur Neuropsychopharmacol* 2003; 13 (3): 199–207.
- 93) Oldham MC, et al. Functional organization of the transcriptome in human brain. *Nat Neurosci* 2008; 11 (11): 1271–1282.
- 94) Ongur D, et al. Glial reduction in the subgenual prefrontal cortex in mood disorders. *Proc Natl Acad Sci U S A* 1998; 95 (22): 13290–13295.
- 95) Ono H, et al. Involvement of thyrotropin in photoperiodic signal transduction in mice. *Proc Natl Acad Sci U S A* 2008; 105 (47): 18238–18242.
- 96) Ozcan ME, et al. Antioxidant enzyme activities and oxidative stress in affective disorders. *Int Clin Psychopharmacol* 2004; 19 (2): 89–95.
- 97) Rajkowska G, et al. Reductions in neuronal and glial density characterize the dorsolateral prefrontal cortex in bipolar disorder. *Biol Psychiatry* 2001; 49 (9): 741–752.
- 98) Ranft K, et al. Evidence for structural abnormalities of the human habenular complex in affective disorders but not in schizophrenia. *Psychol Med* 2010; 40 (4): 557–567.
- 99) Ranjekar PK, et al. Decreased antioxidant enzymes and membrane essential polyunsaturated fatty acids in schizophrenic and bipolar mood disorder patients. *Psychiatry Res* 2003; 121 (2): 109–122.
- 100) Rao AR, et al. Rare deleterious mutations are associated with disease in bipolar disorder families. *Mol Psychiatry* 2017; 22 (7): 1009–1014.
- 101) Rapoport SI. Lithium and the other mood stabilizers effective in bipolar disorder target the rat brain

- arachidonic acid cascade. *ACS Chem Neurosci* 2014; 5 (6): 459–467.
- 102) Rueckert EH, et al. Cis-acting regulation of brain-specific ANK3 gene expression by a genetic variant associated with bipolar disorder. *Mol Psychiatry* 2013; 18 (8): 922–929.
  - 103) Ruzicka WB, et al. Circuit- and diagnosis-specific DNA methylation changes at gamma-aminobutyric acid-related genes in postmortem human hippocampus in schizophrenia and bipolar disorder. *JAMA Psychiatry* 2015; 72 (6): 541–551.
  - 104) Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 2014; 511 (7510): 421–427.
  - 105) Schubert KO, et al. Proteomic pathway analysis of the hippocampus in schizophrenia and bipolar affective disorder implicates 14-3-3 signaling, aryl hydrocarbon receptor signaling, and glucose metabolism: Potential roles in GABAergic interneuron pathology. *Schizophr Res* 2015; 167 (1–3): 64–72.
  - 106) Shamir A, et al. Inositol monophosphatase in immortalized lymphoblastoid cell lines indicates susceptibility to bipolar disorder and response to lithium therapy. *Mol Psychiatry* 1998; 3 (6): 481–482.
  - 107) Shao L, et al. Mitochondrial involvement in psychiatric disorders. *Ann Med* 2008; 40 (4): 281–295.
  - 108) Shiah IS, Yatham LN. Serotonin in mania and in the mechanism of action of mood stabilizers: A review of clinical studies. *Bipolar Disord* 2000; 2 (2): 77–92.
  - 109) Shimizu E, et al. Alterations of serum levels of brain-derived neurotrophic factor (BDNF) in depressed patients with or without antidepressants. *Biol Psychiatry* 2003; 54 (1): 70–75.
  - 110) Siciliano G, et al. Autosomal dominant external ophthalmoplegia and bipolar affective disorder associated with a mutation in the ANT1 gene. *Neuromuscul Disord* 2003; 13 (2): 162–165.
  - 111) So J, et al. Impaired endoplasmic reticulum stress response in B-lymphoblasts from patients with bipolar-I disorder. *Biol Psychiatry* 2007; 62 (2): 141–147.
  - 112) Stelzhammer V, et al. Distinct proteomic profiles in post-mortem pituitary glands from bipolar disorder and major depressive disorder patients. *J Psychiatr Res* 2015; 60: 40–48.
  - 113) Stern S, et al. Neurons derived from patients with bipolar disorder divide into intrinsically different sub-populations of neurons, predicting the patients' responsiveness to lithium. *Mol Psychiatry* 2018; 23 (6): 1453–1465.
  - 114) Steullet P, et al. The thalamic reticular nucleus in schizophrenia and bipolar disorder: role of parvalbumin-expressing neuron networks and oxidative stress. *Mol Psychiatry* 2018; 23 (10): 2057–2065.
  - 115) Suhara T, et al. D1 dopamine receptor binding in mood disorders measured by positron emission tomography. *Psychopharmacology (Berl)*. 1992; 106 (1): 14–18.
  - 116) Sun X, et al. Downregulation in components of the mitochondrial electron transport chain in the post-mortem frontal cortex of subjects with bipolar disorder. *J Psychiatry Neurosci* 2006; 31 (3): 189–196.
  - 117) Suomalainen A, et al. Multiple deletions of mitochondrial DNA in several tissues of a patient with severe retarded depression and familial progressive external ophthalmoplegia. *J Clin Invest* 1992; 90 (1): 61–66.
  - 118) Swift RG, et al. Psychiatric findings in Wolfram syndrome homozygotes. *Lancet (London, England)* 1990; 336 (8716): 667–669.
  - 119) Swift RG, et al. Psychiatric disorders in 36 families with Wolfram syndrome. *Am J Psychiatry* 1991; 148 (6): 775–779.
  - 120) Takaesu Y, et al. Circadian rhythm sleep-wake disorders predict shorter time to relapse of mood episodes in euthymic patients with bipolar disorder: A prospective 48-week study. *J Clin Psychiatry* 2018; 79 (1): 17m11565.
  - 121) Terrillion CE, et al. Reduced levels of Cacna1c attenuate mesolimbic dopamine system function. *Genes Brain Behav* 2017; 16 (5): 495–505.
  - 122) Tkachev D, et al. Oligodendrocyte dysfunction in schizophrenia and bipolar disorder. *Lancet (London, England)* 2003; 362 (9386): 798–805.
  - 123) Tobe BTD, et al. Probing the lithium-response pathway in hiPSCs implicates the phosphoregulatory set-point for a cytoskeletal modulator in bipolar pathogenesis. *Proc Natl Acad Sci U S A* 2017; 114 (22): E4462–E4471.
  - 124) Tomita H, et al. Effect of agonal and postmortem factors on gene expression profile: Quality control in microarray analyses of postmortem human brain. *Biol Psychiatry* 2004; 55 (4): 346–352.
  - 125) Vawter MP, et al. Mitochondrial-related gene expression changes are sensitive to agonal-pH state: Implications for brain disorders. *Mol Psychiatry* 2006; 11 (7): 615, 663–679.

- 126) Williams RS, et al. A common mechanism of action for three mood-stabilizing drugs. *Nature* 2002; 417 (6886): 292–295.
- 127) Xiao Y, et al. The DNA methylome and transcriptome of different brain regions in schizophrenia and bipolar disorder. *PLoS One* 2014; 9 (4): e95875.
- 128) Yang S, et al. Assessment of circadian function in fibroblasts of patients with bipolar disorder. *Mol Psychiatry* 2009; 14: 143–155.
- 129) Yatham LN, et al. Brain serotonin-2 receptors in acute mania. *Br J Psychiatry* 2010; 196 (1): 47–51.
- 130) Yin L, et al. Nuclear receptor Rev-erbalpha is a critical lithium-sensitive component of the circadian clock. *Science* 2006; 311 (5763): 1002–1005.
- 131) Yoon IS, et al. Altered IMPA2 gene expression and calcium homeostasis in bipolar disorder. *Mol Psychiatry* 2001; 6 (6): 678–683.
- 132) Yoshimi N, et al. Cerebrospinal fluid metabolomics identifies a key role of isocitrate dehydrogenase in bipolar disorder: Evidence in support of mitochondrial dysfunction hypothesis. *Mol Psychiatry* 2016; 21 (11): 1504–1510.
- 133) Yurgelun-Todd DA, et al. fMRI during affect discrimination in bipolar affective disorder. *Bipolar Disord* 2000; 2 (3 Pt 2): 237–248.
- 134) Zhao Z, et al. Transcriptome sequencing and genome-wide association analyses reveal lysosomal function and actin cytoskeleton remodeling in schizophrenia and bipolar disorder. *Mol Psychiatry* 2015; 20 (5): 563–572.